

Physiological and morphometric characteristics of *Pachycereus pringlei* (S. Watson) Britton & Rose seedlings applying organic manures

Joselyn Vanessa Seminario-Peña¹, Alejandra Nieto-Garibay^{1*}, Enrique Troyo-Diéguez¹, Bernardo Murillo-Amador¹, Alfonso Medel-Narváez¹, Teresa Terrazas²

¹Centro de Investigaciones Biológicas del Noroeste, S.C. La Paz, Baja California Sur, México.

²Instituto de Biología de la Universidad Nacional Autónoma de México, Ciudad de México, México

*Corresponding author: anieto04@cibnor.mx

Abstract. The establishment of *Pachycereus pringlei* seedlings is a scarce event; soil conditions are one of the factors attributed to its high mortality rate in the early stages of development. The use of organic manures as a substrate helps to improve the structure, porosity, and density of the soil where they live, in addition to providing nutrients that would benefit the roots, increasing the possibility of their establishment during the emergence stage. The objective of this study was to evaluate the effect of different proportions of organic manures as a sown substrate in the emergence and establishment of *Pachycereus pringlei* seedlings, using nine treatments of organic manures in a completely randomized experimental design with four replications. The variables measured were emergence rate (ER) and percentage (EP) and their indices, seedlings' morphometric, and physiological characteristics in the establishment stage. The results showed that the evaluated variables in the emergence and seedling establishment phase indicate the preference for natural substrate and organic manure combinations in proportions of 50 and 30%, respectively. The analysis of variance showed significant differences regarding the morphometric and physiological variables of the stem and root of the evaluated treatments. The emergence and establishment analyses show that carbon seedlings under different organic manure treatments and controlled conditions had excellent survival results compared to the low rates of recruitment and survival reported under natural conditions. Similarly, the implementation of organic manures improved the physiological and morphometric characteristics of the carbon seedlings compared to the control treatment with the natural substrate.

Citation: Seminario-Peña, J.V., Nieto-Garibay, A., Troyo-Diéguez, E., Murillo-Amador, B., Medel-Narváez, A. and Terrazas, T. 2023. Physiological and morphometric characteristics of *Pachycereus pringlei* (S. Watson) Britton & Rose seedlings applying organic manures. *Journal of the Professional Association for Cactus Development*. 25: 36-58. <https://doi.org/10.56890/jpacd.v25i.516>

Associate Editor: Pablo Preciado-Rangel

Technical Editor: Tomas Rivas-Garcia

Received date: 13 January 2023

Accepted date: 03 April 2023

Published date: 22 May 2023



Copyright: © 2023 by the authors.

Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY NC SA) license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>).

Keywords: emergence, establishment, cacti, compost, vermicompost

Introduction

The Cacti from arid and semi-arid zones where water is limited have unique morphometric, physiological, and anatomical characteristics that allow them to tolerate extreme climatic conditions and complete their life cycle (Nobel *et al.*, 1988; Bravo-Hollis and Scheinvar, 1995). These characteristics for adapting to environments include crassulacean acid metabolism (CAM), thick cuticle, and leaves modified to spines (Granados *et al.*, 1998). Cacti are vulnerable species in the early stages of development (Martínez-Ramos *et al.*, 2016) because of abiotic and biotic factors (Turner, 1990; Valiente-Banuet and Ezcurra, 1991; Mandujano *et al.*, 1997). Cacti seedlings must grow as fast as possible, establish roots for rapid water absorption, and compete for nutrients, light, and space with other plants (Fenner and Thompson, 2005). The most important source of nutrients for plants is organic matter, which in arid regions is limited; therefore, many cacti seedlings do not reach the next stage and die due to a lack of nutrients (Ramirez, 2011).

The early stages of cacti are considered critical, as the seedlings depend on environmental variables for their survival and establishment, such as moisture, light, and nutrients (Sánchez-Soto et al., 2010; Pimienta-Barrios et al., 2012). The emergence is the process where the appearance of the seedling is observed on the soil surface; it is a stage after the seed germination or sprouting of the bud. On the other hand, the seedling establishment is defined as the process by which a plant establishes itself, resists, and survives. It also includes the stages of germination and emergence until the development of true leaves -spines in cacti- (National Research Council, 2002; Dumroese et al., 2012).

The soil represents an important factor in the first stages of plant development, which is why in recent years, the use of organic manures (compost and vermicompost) has increased due to the benefits it has, among which are the increased permeability and retention of moisture in the soil for longer, has a buffer capacity against sudden changes in acidity, and another advantage is the supply of nutrients, which improves the morphometric and physiological characteristics of plants (Bashan et al., 2009; Nieto-Garibay et al. 2021). Some morphometric and physiological changes that can be evidenced depending on the environment in which these species are found are the presence of a thick cuticle, high osmotic concentrations to extract moisture from very dry soils, great development of the root system, and size of the portion reduced area (González, 2012).

Pachycereus pringlei (S.Watson) Britton & Rose is a species of cactus endemic to the Sonoran Desert, which is a reservoir of nutrients, provides shelter and is a source of food for local fauna (Delgado-Fernández et al., 2017). This species has a high germination rate (Suzán-Azpíri and Sosa, 2006; Bacilio et al., 2011); however, the establishment of seedlings is a sporadic event due to high mortality rates (Bullock et al., 2005), which could be linked to the edaphic conditions of arid environments, such as the low percentage of organic matter (Pérez-Sánchez et al., 2015). This study aimed to evaluate the effect of different proportions of organic manures as a sown substrate in the emergence and establishment of *Pachycereus pringlei* seedlings.

Material and Methods

Study site

The experiments of emergence and establishment were carried out in the Center for Biological Research of Northwest México S.C (CIBNOR), west of La Paz city, Baja California Sur (BCS), México located at 24°08'10.03" N and 110°25'35.31" W.

Fruits collection

The ripe cardon fruits were collected between June to July at the Biological Station "Dra. Laura Arriaga-Cabrera" located at 24° 07'41.4" N and 110° 26' 19.0" W (Figure 1a). The seeds were extracted from the fruit through longitudinal dissections and later were dried at room temperature using paper napkins and cotton (Figure 1b).




Figure 1. (a) *Pachycereus pringlei* fruits collected between June and July; (b): Seeds extraction from the fruits and drying at room temperature.

Experimental design

The experiments of the emergence and establishment of seedlings stages were carried out using a completely randomized experimental design where the main study factor was the sowing substrate with nine treatments and four replications of 50 seeds. The treatments consisted in the combination of the substrate in different proportions of organic manure and native soil (Table 1). The organic manures were from agricultural residues and obtained from Rancho Buena Suerte (organic farm) in the town of La Paz, BCS. The physicochemical composition, electrical conductivity (EC), organic matter (OM), apparent density (AD), porosity (Po), and pH of the treatments were analyzed in the soil science laboratory of CIBNOR (Table 1). The substrate's nutrient content (NH_3 , NO_3^- , SO_4^{2-} , Ca, P, Mg, and K) were analyzed using a portable photometer (9, Hanna Instruments Inc., Woonsocket, RI, USA).

1 **Table 1.** Physicochemical composition of each treatment (sowing substrates).

Treatments	EC dS m ⁻¹	OM %	PO %	DA G cm ⁻³	pH	NO ₃ ⁻ mg L ⁻¹	NH ₃	SO ₄ ²⁻	Mg	P	Ca	K
NS100 (control)	0.4	0.4	19.25	1.51	7.45	ND	0.006	0.1	0.5	0.008	0.9	0.1
C100	30.6	39.6	22.26	0.41	8.33	2.17	0.04	16.7	0.3	0.44	156.7	11.0
V100	15.2	20.8	10.19	0.73	8.16	1.17	0.12	16.7	21.7	0.60	176.7	96.7
C30NS70	6.8	14.6	1.89	1.05	8.42	0.67	0.36	13.3	5.0	0.40	166.7	8.7
C70NS30	21.3	30.1	17.36	0.54	8.79	1.33	0.13	20.0	6.7	0.27	116.7	10.5
C50NS50	10.6	16.7	4.91	0.87	8.51	0.83	0.08	15.0	13.3	0.47	110.0	9.5
V30NS70	5.2	3.1	15.85	1.42	8.50	1.00	0.43	18.3	5.0	0.43	200.0	11.7
V70NS30	8.7	18.6	3.02	1.08	8.45	1.17	0.14	15.0	6.7	0.50	160.0	8.7
V50NS50	9.1	6.3	9.96	1.24	8.38	0.50	0.10	15.0	5.0	0.57	166.7	10.5

2 NS100: 100% Native Soil. C100: 100% Compost. V100: 100% Vermicompost. C30NS70: Compost 30%-Native Soil 70%. C70NS30: Compost
3 70%-Native Soil 30%. C50NS50: Compost 50%-Native Soil 50%. V30NS70: Vermicompost 30%-Native Soil 70%. V70NS30: Vermicompost
4 70%-Native Soil 30%. V50NS50: Vermicompost 50%-Native Soil 50%. EC: electrical conductivity. OM: Organic matter. PO: Porosity. AD:
5 Apparent density. ND: Not detected. (*n* = 3).

6 Seedlings emergence stage

7 The seeds were sown in trays of 200 cavities divided previously into four sections of 50 cavities,
8 each representing one replication of 50 seeds. The trays were covered with plastic micro-
9 tunnels to prevent rapid evaporation and placed inside a 2.5 × 2.5 m anti-aphid mesh. The
10 treatments were irrigated every 4 to 7 days with water from the desalination plant of the
11 experimental field. The emergence was considered when seedlings began to sprout 2 mm
12 above the substrate and were recorded daily until 33 days after sowing.

13 Emergence Indices

14 Through the emergence indices, the favorable performance of seedling emergence was
15 determined. Table 2 describes the emergence indices calculated using the “GerminaR”
16 package in RStudio (Lozano-Isla et al., 2018).

17 **Table 2.** Description of emergence indices.

Emergence Indices	Units	Formulas	Reference
Emergence Percentage (EP)	%	$EP = \frac{n_i}{N} \times 100$ Where: n_i = Seedlings emerged on the i^{th} day N = Total number of seedlings in the experiment	Maguire (1962)
Speed of emergence (SE)	%	$SE = \sum_{i=1}^k \frac{n_i}{n_i X_i} \times 100$ Where: n_i = Seedlings emerged on the i^{th} day X_i = Number of days since seed sowing k = The last day of emergence evaluation	Ranal and Santana (2006)
Emergence Index (EI) or Mean Emergence Time	time	$EI = \sum_{i=1}^k \frac{n_i t_i}{n_i} = \frac{1}{MER}$ Where: n_i = Seedlings emerged on the i^{th} day t_i = Time from the start of the experiment to the i^{th} day of observation k = The last day of emergence evaluation MER = Mean emergence rate	Ranal and Santana (2006)
Emergency Rate Index (ERI)	time	$ERI = \frac{\sum_{i=1}^k n_i}{MER}$ Where: n_i = Seedlings emerged on the i^{th} day k = The last day of emergence evaluation MER = Mean emergence rate	Bilbro and Wanjura (1982)

Mean Emergence Rate (MER)	time ⁻¹	Where: EI = Emergence Index	$MER = \frac{1}{EI}$	Ranal and Santana (2006)
----------------------------------	--------------------	--------------------------------	----------------------	--------------------------

20

21

Seedlings establishment stage

23 The seedlings' establishment stage was evaluated at the end of the emergence stage and until
24 153 days after sowing. The physiological and morphometric measurements of survived
25 seedlings were evaluated individually in the plant physiology laboratory of CIBNOR, selecting
26 three seedlings per replication and treatment. The treatments were irrigated every 10 days with
27 water from the desalination plant of the experimental field during this phase.

28

Survival curve

30 The seedlings' establishment was performed using Kaplan-Meier survival curves (Kaplan and
31 Meier, 1958) and compared using the Mantel-Haenszel log-rank analysis (Mantel and
32 Haenszel, 1959). Survival curves were evaluated (Figure 2) from the day 33 when an
33 asymptote was observed in the emergence of seedlings, and 5 treatments that registered at
34 least 1 dead individual were included. The treatments C30NS70, C70NS30, V30NS70, and
35 V70NS30 were discarded from the analysis because they did not register the deaths of
36 individuals during the evaluated period.

37

$$38 \quad S(t_i) = \prod_{t_i \leq t} \frac{n_i - d_i}{n_i}$$

39

40 Where, n_i is the number of living individuals before the time t_i , d_i is the number of events at
41 the time t_i , and t_i is the inter-occurrence time of the i^{th} event in the i^{th} research unit.

42

Morphometric variables (stems and roots)

44 The spine length, the primary root length, diameter, and stem length were measured using a
45 digital vernier (Cole-Parmer, General model 143). The number of spines was determined for
46 each seedling.

47

48 The dry and fresh weight of stems and roots were determined using an analytical balance
49 (Mettler Toledo, model AG204). The stems and roots were placed in paper bags and placed in
50 a laminar flow oven (Sheb-Lab, model FX-5, series 1000203) for 24 hours at 70°C until entirely
51 dehydrated to obtain the dry weight.

52

53 The root area was determined using an area meter (LICOR, model LI-1300) scanning the entire
54 root. Previously, the roots were rinsed with topwater for 5 min to remove the excess substrate
55 and stored in the refrigerator until scanning. The values for the root area variable are expressed
56 in centimeters squared (cm²).

57

Physiological variables

58

59 *Relative water content (RWC)*. The RWC was calculated using the formula proposed by
60 Yamasaki and Dillenburg (1999):

$$61 \quad RWC = \frac{F_w - T_w}{D_w - T_w} \times 100$$

62
63 Where, F_w is the fresh weight, D_w is the dry weight, and T_w is the turgent weight. The spines of
64 the stems were removed and subsequently weighed to obtain the fresh weight (F_w). To
65 determine the turgid weight (T_w), stems were placed in distilled water inside a closed Petri dish
66 for 24 hours. After that, the sample of the stems was placed in a laminar flow oven (Sheb-Lab,
67 model FX-5, series 1000203) at 70°C for 24 hours to obtain the dry weight (D_w). The weights
68 of the plant material were obtained using an analytical balance (Mettler Toledo, model AG 204).
69 *Osmotic potential (Ψ_o)*. The measurements of osmotic potential were made according to the
70 method proposed by Moghaieb et al. (2004). The seedlings were placed in vials and
71 subsequently placed in an ultra-freezer at -20 °C. The vials were thawed and centrifuged at
72 1200 rpm x g for 20 min at 4°C to extract the cells from the sap. The osmotic potential was
73 determined with a vapor pressure osmometer (Wescor, VAPRO model). The data were
74 transformed to MPa using the formula proposed by Van't Hoff (Salisbury and Ross, 1992),
75 which states the following:

$$76 \quad \Psi_o = -C \times R \times T$$

77
78
79 Where, C is the molarity of the solution (mol of the solute kg⁻¹ H₂O), R is the universal gas
80 constant (0.0831 kg Mpa mol⁻¹ K⁻¹), and T is the absolute temperature (K).

81
82 *Water potential (Ψ_w)*. The measurements of water potential were made at the critical time
83 (approximately 12:00 h), considering the interval in which the highest temperatures of the day
84 occur. The water potential was determined using a dew point potentiometer (Dewpoint
85 PotentiaMeter, model WP4-T). Three readings of each treatment were made, using complete
86 samples of cardon seedlings at the end of the experiment.

87
88 *Turgor pressure (Ψ_t)*. The turgor pressure was calculated considering the difference between
89 the water potential (Ψ_w) and the osmotic potential (Ψ_o) (Moghaieb et al., 2004).

$$90 \quad \Psi_t = \Psi_w - \Psi_o$$

91
92
93 *Chlorophyll a, b, and total*. A hollow steel punch was used to obtain the fresh tissue discs, to
94 which 80% acetone was added, and it was left to stand for 48 hours in a test tube at room
95 temperature and in the dark. Subsequently, the absorbance was read in a UV/Visible
96 spectrophotometer (HACH, model DR 3900). Chlorophyll contents were calculated with the
97 equation proposed by Strain and Svec (1996):

$$98 \quad \text{Chl } a \text{ (mg mL}^{-1}\text{)} = 11.64 \times (A663) - 2.16 \times (A645)$$

$$99 \quad \text{Chl } b \text{ (mg mL}^{-1}\text{)} = 20.97 \times (A645) - 3.94 \times (A663)$$

100
101

102 Where, A_{663} and A_{645} represent the absorbance values read at a wavelength of 663 nm and
103 645 nm, respectively. Additionally, total chlorophyll was measured indirectly, through the sum
104 of chlorophylls a and b, to obtain a prediction model of total chlorophyll values.

105

106 **Statistical analyses**

107 The normality and homoscedasticity of variances of the data were verified, and subsequently,
108 a one-way ANOVA test was performed. The mean differences among treatments of each
109 variable were detected with a mean difference test (Tukey HSD, $p \leq 0.05$). A nonparametric
110 Kruskal-Wallis test was performed when the data did not meet the statistical assumptions,
111 followed by Dunn's post-hoc test for group comparisons. Statistical analysis was performed
112 with the free access software RStudio. The significance levels for each treatment were later
113 identified using the "Agricolae" and "MultcompView" packages (De Mendiburu, 2014; Graves
114 *et al.* 2019).

115

116

117

117 **Results and discussion**

118

119 **Seedlings emergence stage**

120 *Emergence Indices*

121 The results showed statistically significant differences ($p < 0.05$) in the emergence indices
122 evaluated for each treatment in this study (Table 3). Regarding the percentage of emergence
123 (EP), the highest values were obtained for the treatments V30NS70, C30NS70, V50NS50, and
124 C50NS50. On the other hand, the 100% treatments with organic manures registered low values
125 of the emergence percentage (EP) and slow emergence speed (SE). Treatments NS100 and
126 C30NS70, which showed the first seedlings emerged from the substrate, registered the highest
127 emergence speed (SE) values. Concerning the emergence index (EI), which estimates the time
128 it takes for the seedling to emerge, the highest values were recorded for the V100 and C100
129 treatments because the mean total seedling emergence time for these treatments was slower
130 than the other treatments. The treatments with an organic substrate, including the control
131 (NS100), that was watered every 4 to 7 days, showed more than 19 % of the seedlings'
132 emergence. These results are similar to those reported by Puente (2004), who described that
133 in the control treatment with only saturation irrigation, cardon seeds had a high germination
134 percentage. Marchiol *et al.* (1999) subjected different types of substrate treatments with
135 compost to seeds from herbaceous plants and found that the lowest germination percentage
136 was recorded for the 100 % compost treatment. These results are similar to the treatments of
137 C100 and V100 of this study, which registered less than 34 % of emerged seedlings. In addition,
138 there was a delay in the emergence phase, evidenced by the speed of emergence (SE) and
139 the emergence index (EI), which could be associated with the combination of high levels of
140 electrical conductivity and potassium, which affects germination in some cactus (Teixeira *et al.*,
141 2004; Sarria-Perea, 2010; Freire *et al.*, 2018).

142

143 The emergence percentage increased by more than 25% for the V30NS70 treatment
144 concerning control group. Organic manures in proportions of 50 and 30 % for C30NS70,
145 V50NS50, and C50NS50 also presented a high percentage (>58%) of seedlings emerged on
146 day 33. Likewise, the seedling emergence variables showed an important variation concerning

147 the C100 and V100 treatments, which indicates the preference for the combination of natural
148 substrate and organic manure in proportions of 50% and 30 %, which is corroborated with the
149 emergence indices (Table 3).

150

151 The NS100 and C30NS70 treatments recorded the highest values of emergence speed (SE)
152 and emergence rate index (ERI) since on days 1 and 2 the first seedlings began to sprout from
153 the substrate. The average speed of emergence for *P. pringlei* in the treatments with organic
154 manures evaluated was 8 days. Regarding the above, Godínez-Alvarez and Valiente-Banuet
155 (1998) mention that the seeds of cacti species germinate quickly in the first weeks when they
156 have moisture, for example, some species of *Opuntia* take to germinate an average of 13 days,
157 but under controlled conditions, they germinate between 5 and 6 days (Gonzalez-Cortés *et al.*,
158 2018).

159

160 **Table 3.** Emergence indices of cardon seedlings under organic manure treatments.

Treatments	EP	SE	EI	ERI	MER
	%	%time ⁻¹	time ⁻¹	time ⁻¹	time
NS100 (control)	57.5 ± 10.6 ^{abc}	12.839 ± 0.73 ^a	7.806 ± 0.42 ^c	3.683 ± 0.731 ^{ab}	0.1283 ± 0.0073 ^a
C100	19 ± 2.58 ^d	6.129 ± 0.29 ^c	13.315 ± 0.78 ^a	0.581 ± 0.064 ^c	0.0612 ± 0.0029 ^c
V100	33 ± 6.83 ^{cd}	5.876 ± 0.29 ^c	17.050 ± 0.83 ^a	0.968 ± 0.192 ^c	0.0587 ± 0.0029 ^c
C30NS70	69.5 ± 12.4 ^{ab}	10.799 ± 1.26 ^{ab}	9.347 ± 0.99 ^{bc}	3.718 ± 0.429 ^{ab}	0.1079 ± 0.0126 ^{ab}
C70NS30	54 ± 16.6 ^{abc}	8.144 ± 0.99 ^{bc}	12.415 ± 1.48 ^{ab}	2.175 ± 0.707 ^{bc}	0.0814 ± 0.0099 ^{bc}
C50NS50	59 ± 9.31 ^{abc}	9.874 ± 2.29 ^{ab}	10.618 ± 2.84 ^{bc}	2.778 ± 0.467 ^{ab}	0.0987 ± 0.0229 ^{ab}
V30NS70	83 ± 22.5 ^a	10.429 ± 2.17 ^{ab}	9.993 ± 2.60 ^{bc}	4.153 ± 1.810 ^a	0.0104 ± 0.0217 ^{ab}
V70NS30	47 ± 4.76 ^{bcd}	9.094 ± 1.22 ^{bc}	11.148 ± 1.51 ^{ab}	2.108 ± 0.380 ^{bc}	0.0909 ± 0.0122 ^{bc}
V50NS50	66 ± 14 ^{ab}	10.453 ± 1.22 ^{ab}	9.666 ± 1.13 ^{bc}	3.414 ± 0.613 ^{ab}	0.1045 ± 0.0229 ^{ab}

161 NS100 (control): 100% Native Soil. C100: 100% Compost. V100: 100% Vermicompost. C30NS70: Compost 30%-Native Soil 70%. C70NS30: Compost 70%-Native
 162 Soil 30%. C50NS50: Compost 50%-Native Soil 50%. V30NS70: Vermicompost 30%-Native Soil 70%. V70NS30: Vermicompost 70%-Native Soil 30%. V50NS50:
 163 Vermicompost 50%-Native Soil 50%. EP: Emergence percentage. SE: Speed of emergence. EI: Emergence Index. ERI: Emergency rate index. MER: Mean
 164 emergence rate. Means ± standard deviation followed by the same letter in the same column are not significantly different. The significance level was set at $p \leq 0.05$.
 165 (n=4).
 166

167 **Seedlings establishment stage**

168 *Survival curve*

169 No significant differences were found in seedling survival during the seedling establishment stage between treatments of organic
 170 manures and control (LR=7.02, g.l.=4, $p > 0.05$). The establishment showed that cardon seedlings had high survival, with 1 to 2 deaths
 171 recorded after 153 days (Figure 2), compared to the low rates of recruitment and survival recorded under natural conditions (Bullock *et*
 172 *al.*, 2005). These results are related to the availability of nutrients and the retention of humidity as benefits of the use of organic manures.
 173 It is noteworthy that the survival success of seedlings, and especially cacti, can be determined by the amount of water stored in the
 174 tissues during the first stage of growth, which is favored by water retention by vermicompost and compost.




Figure 2. Survival of cardon seedlings under organic manure treatments.

The arid and semi-arid zones are characterized by environments where water resources are limited, and temperatures are extreme. These zones have soils with low nutrient availability and atmospheric decompensation due to water loss by evapotranspiration (Vásquez-Méndez *et al.*, 2011; Montañaño *et al.*, 2016). These characteristics cause cacti seedlings vulnerable in the early stages of life, with slow growth rates (Martínez-Ramos *et al.*, 2016).

Morphometric variables of the stems

The analysis of variance showed significant differences ($p < 0.05$) with respect to the variables evaluated of the stem between treatments of organic manures and control (Figure 3). It is observed that the treatment of NS100 (control) registers the thinnest and smallest stems, unlike the treatments with organic manures, which retain a higher humidity of the substrate. It is observed that the largest seedlings were registered for the treatment of C30NS70, C50NS50, C70NS30, V30NS70, V50NS50 and V70NS30. Even though the C100 and V100 treatments registered low emergence percentages, the few seedlings that emerged were larger than the NS100 treatment (control). The morphometric variables of the stem showed that the combination of organic manures and native substrate allowed rapid growth of the stems, due to the contribution of nutrients and moisture retention by the organic substrate (Domínguez, 2004; Doan *et al.*, 2015; Nieto-Garibay *et al.*, 2021). The smallest seedlings were recorded the treatments with extreme values of electrical conductivity and organic matter, the NS100 treatment had the lowest values of these parameters, while the V100 and C100 treatments were the opposite (Figure 3). According to González and Chueca (2010), Celaya-Michel and Castellanos-Villegas, (2011), the high saline content of soils can affect the development of the seedlings leading to water deficit; this can be a consequence of the use of pure proportions of organic manures which might hamper the growth in seedlings. The proportions combined with organic manures showed better seedling development than to the control treatment (NS100) (Figure 3). The above is supported by Uddin *et al.*, 2012, who evidenced a positive effect application of organic manure with soil in leguminous seedlings.

Likewise, it is observed that there were significant differences ($p < 0.5$) in the fresh weight of the seedlings between treatments. Graphs 3a and 3b show a directly proportional relationship between the stems' size and the fresh weight of each treatment. The number of spines also showed statistically significant differences ($p < 0.5$) between the control (NS100) and organic manure treatments. The combined treatments of organic manures showed a more substantial number of spines than the control group (NS100) and the pure treatments with organic manures (V100 y C100). Likewise, while counting the spines, it was possible to observe that the combined treatments

had thicker spines while the control treatment had thinner spines. The carbon seedlings exhibited morphometric characteristics like those registered for other species of the Pachycereeae tribe, where it was observed as they grew that the first areoles had tiny spines and whitish trichomes (Loza-Cornejo and Terrazas, 2011).




Figure 3. Average values (\pm SD) of stem morphometric variables of cardon seedlings under organic manure treatments, (a) Stem diameter; (b) Stem length; (c) Number of spines; (d) Stem fresh weight. The bars followed by the same letter are not significantly different (Tukey HSD $p \leq 0.05$). The vertical lines at the top of the bars represent the standard deviation.

Morphometric variables of the roots

There are statistically significant differences ($p < 0.5$) in the evaluated variables of the root between treatments of organic manures and control (Figure 4). The same trend is observed with the treatments' seedlings of C30NS70, C50NS50, C70NS30, V30NS70, V50NS50, and V70NS30, which acquired seedlings of greater diameter and length showing higher root area values. Similarly, the same pattern is repeated for the average length of the tap root. The treatments of C100, V100, and NS100 showed little root growth, which may be related to the scarcity and ratio of organic manure used for these treatments as indicated previously. Loza-Cornejo *et al.* (2003) mention that the success of the establishment of cacti depends on the root system responsible for the absorption of water and nutrients from the soil. There is a relationship between the area of the root and the aerial part of the seedling since the treatments of NS100, V100, and C100 reported the thinnest and smallest stems. That is, as the plant has more root hairs, the size of the plant should be due to the greater absorption of water and nutrients (Niklas *et al.*, 2000, 2002).




Figure 4. Average values (\pm SD) of root morphometric variables of carbon seedlings under organic manure treatments, (a) Root area; (b) Root length; (c) Root fresh weight; (d) Root dry weight. Significance levels were given by the Tukey test. The bars with different letters indicate significant differences at $p < 0.05$, and the vertical lines at the top of the bars represent the standard deviation.

Physiological variables

Relative water content

In arid zones, the ability of a plant to maintain its water status as the soil water potential decreases constitutes an adaptation to water deficit, which is reflected in the relative water content (RWC) (Ortiz *et al.*, 2003). It can be seen in Figure 5a, that the seedlings of the treatments do not present significant differences concerning the relative water content. Considering the importance of water in plants and the adaptations that cacti have developed to use water efficiently and to be able to survive in adverse conditions, it can be considered essential to know the water status of CAM plants. Even though the seedlings of the V100, V50NS50, and C100 treatments register shallow values of water potential (Figure 5c), it can be observed that the RWC was greater than 80% for these treatments. Cacti tissues have large vacuoles that occupy more than 90% of the cell volume, which improves their water storage capacity (Pérez-Sánchez *et al.*, 2015). This ability could explain why most treatments' RWC values are not as marked, and according to Flexas and Medrano (2002), the seedlings evaluated from 9 treatments showed moderate (70-85%) to slight (85-95%) stress.

Osmotic potential

The variable osmotic potential or osmotic pressure is expressed in megapascals (MPa) and is the concentration of osmotically active solutes (soluble sugars, ions, and amino acids) in the vacuole (Taiz and Zeiger, 2006; Salmon *et al.*, 2020), which indicates the water status of the plant (Azcón-Bieto and Talón, 2013). The results showed that osmotic potential was affected in the C100, V100, and NS100 treatments (Figure 5b), registering the most negative values and smallest seedlings concerning the treatments with organic manures in proportions of 50 and 30 %. The above indicates that the stems under these treatments have dehydrated cells due to the low concentration of malate in the vacuoles (Azcón-Bieto y Talón, 2013; Delgado-Sánchez *et al.*, 2017). However, cacti can adjust osmotically to tolerate dry environments (Goldstein *et al.*, 1991; Delgado-Sánchez *et al.*, 2013), and this can be compared in the present study, where the measured values of osmotic potential at the level of the lowest ones do not exceed < -1.2 MPa even though during the establishment phase it was irrigated every 10 days.

Water potential

The results show that the V100, C100 and NS100 treatments, where the irrigation water evaporated faster, recorded the most negative values of water potential and smaller seedlings compared to the other treatments. According to the Hsiao (1973) classification of water potential levels, the seedlings subjected to the treatment C30NS70 registered mild stress, moderate stress under C70NS30, C50NS50, V30NS70, and V70NS30 and severe stress with V50NS50, C100, V100, and NS100 (Figure 5c). Agentel *et al.* (2006) mentioned that the RWC and water potential influence the water relationships of plants. Likewise, the water deficit not only occurs when there is little water available in the environment, but also due to low temperatures and high soil salinity (Moreno, 2009). As it was explained in previous points, the treatments that registered severe stress have high levels of electrical conductivity, which cause the water potential of the seedlings more negative. However, cacti have developed adaptations and strategies that allow them to survive in these limiting conditions. Nilsen and Orcutt (2000) mention that xerophytic plants adapt to arid environments and can tolerate water potentials more negative than -4 MPa, which coincides with the results of this study related to the water potential and survival.

Turgor pressure

The variance analysis of the turgor pressure variable showed statistically significant differences ($p < 0.5$) between the organic manures and control treatments (Figure 5d). The most negative turgor pressure treatment was recorded for V100. The turgor pressure suggests that the V100, NS100, V50NS50, and C100 treatments have less rigid cell walls, which would be related to the high content of EC in the pure proportions of organic manures that causes dehydration in the cells (Nobel, 2006; Schuch and Kelly, 2008). Furthermore, the control treatment without organic substrate was also affected by the turgor loss due to the irrigation water's rapid volatilization.




Figure 5. Average values (\pm SD) of physiological variables of cactus seedlings under organic manure treatments, (a) Relative water content; (b) Osmotic potential; (c) Water potential; (d) Turgor pressure. Significance levels were given by the Tukey and Dunn test. Different letters above the bars indicate significant differences at $p < 0.05$, and the vertical lines at the top of the bars represent the standard deviation.

Chlorophyll a, b, and total

The content of chlorophyll a, b and total did not show significant differences between treatments (Figure 6); however, the highest average values of total chlorophyll were recorded for the C70NS30, V30NS70, and V100 treatments this pattern was recorded in the same way for chlorophyll a and chlorophyll b. The treatments with lower values of chlorophyll a, b, and total were recorded for C100 and NS100 (control). The total chlorophyll content with the organic manure treatments increased in 29 a 40 $\mu\text{g cm}^{-2}$. In studies using organic manures, there has been an increase in chlorophyll content which is directly related to good nutritional status (Degli-Esposti *et al.*, 2011; Flores *et al.*, 2022), while in substrates where the salt content is high, the presence of photosynthetic pigments decreases due to the inhibition of specific enzymes responsible for chlorophyll synthesis (Franco-Salazar and Véliz, 2008; Naseer *et al.*, 2022). These results are similar to those reported by Meléndez *et al.* (2006) and Trujillo *et al.* (2010) who mention that the chlorophyll content decreases in soils with low moisture gradients and increases in soils with high moisture gradients. On the other hand, stress due to water deficit can also produce photooxidation and degradation of photosynthetic pigments, such as chlorophyll (Flexas and Medrano, 2002; Anjum *et al.*, 2011).




Figure 6. Average values (\pm SD) of physiological variables of cardon seedlings under organic manure treatments, (a) Chlorophyll a; (b) Chlorophyll b; (c) Total chlorophyll content. Significance levels were given by the Tukey and Dunn test. Different letters above the bars indicate significant differences at $p < 0.05$, and the vertical lines at the top of the bars represent the standard deviation.

Conclusions

The results reinforce the information about the benefits by the organic manures of *P. pringlei* during the emergency and establishment phase. Incorporating organic manures in proportions of 50 and 30 % recorded higher values in different emergency indexes and increased the emergence of cardon seedlings. However, the pure treatments with organic manures decreased the emergence of cardon seedlings due to the combination of high levels of electrical conductivity and potassium content.

The establishment of seedlings was high in all treatments. Nevertheless, the seedlings under treatments with proportions of 30, 50 and 70 % of organic manures recorded higher values in the morphometrics variables of the stem and root, which is attributed to the retention of moisture and the supply of nutrients by organic manures, which are essential factors in the first stage of their life cycle. The physiological variables corroborated that cardon seedlings can tolerate negative water potentials of up to -4MPa ; it was also possible to observe that the pure proportions of organic manures registered low levels of osmotic potential. The total chlorophyll content was not affected by using different substrates; however, there was a tendency to favor organic manure treatments.

Considering the importance of water in plants and the adaptations that cacti have developed to make this resource more efficient and to survive in adverse conditions, it is essential to know the water status of CAM plants and strategies to ensure establishment in the early stages of life.

Organic manures are a hopeful tool to increase the emergence and establishment of seedlings of *Pachycereus pringlei* (S.Watson) Britton & Rose and improve their morphological and physiological characteristics. In the same way, these results could have important implications for planning reforestation practices for cacti in danger of extinction.

Ethics statement

Not applicable

Consent for publication

Not applicable

Availability of supporting data

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study is part of the doctoral research of Joselyn Vanessa Seminarío Peña at CIBNOR; with fellowship Conacyt N° 937443. Funding for this research was provided by Conacyt-Coscyt 2020 through the project “Manejo de residuos orgánicos a través del compostaje”.

Author contributions

Conceptualization, J.S., A.N.G. and A.M.N.; methodology, J.S., A.N.G., T.T., A.M.N., E.T.D. and B.M.A.; formal analysis, J.S. and E.T.D.; investigation, J.S.; resources, A.N.G.; data curation, J.S. and E.T.D.; writing—original draft preparation, J.S. and A.N.G.; writing—review and editing, J.S., A.N.G., T.T. and B.M.A.; visualization, A.N.G.; supervision, A.N.G.; project administration, A.N.G.; funding acquisition, A.N.G.

Acknowledgments

The authors are grateful to the technicians Maria Del Carmen Mercado Guido and Lidia Hiraes Lucero of the Plant Physiology Laboratory of CIBNOR, the field Technicians José Raymundo Ceseña Núñez, Pedro Luna García, Adrián Jordán Castro, and Saúl Edel Briseño Ruiz provided valuable aid, and Alvaro Gonzalez Michel of the Hydrology and Irrigation Laboratory of CIBNOR.

References

- Anjum, S.A., Wang, L., Farooq, M., Khan, I. and Xue, L. 2011. Methyl jasmonate-induced alteration in lipid peroxidation, antioxidative defence system and yield in soybean under drought. *Journal of Agronomy and Crop Science*. 197(4): 296–301. <https://doi.org/10.1111/j.1439-037X.2011.00468.x>
- Argentel, L., González, L.M., Ávila, C. and Aguilera, R. 2006. Comportamiento del contenido relativo de agua y la concentración de pigmentos fotosintéticos de variedades de trigo cultivadas en condiciones de salinidad. *Cultivos Tropicales*. 27(3): 49–53.
- Azcón-Bieto, J., and Talón, M. 2013. Fundamentos de fisiología vegetal. McGraw-Hill Interamericana de España, S.L., España. 669 p.

- Bacilio, M., Vazquez, P. and Bashan, Y. 2011. Water versus spacing: A possible growth preference among young individuals of the giant cardon cactus of the Baja California Peninsula. *Environmental and Experimental Botany*. 70(1): 29–36. <https://doi.org/10.1016/j.envexpbot.2010.06.004>
- Bashan, Y., Salazar, B., Puente, M.E., Bacilio, M. and Linderman, R. 2009. Enhanced establishment and growth of giant cardon cactus in an eroded field in the Sonoran Desert using native legume trees as nurses plants aided by plant growth-promoting microorganisms and compost. *Biology and Fertility of Soils*. 45(6): 585–594. <https://doi.org/10.1007/s00374-009-0367-x>
- Bilbro, J.D. and Wanjura, D.F. 1982. Soil crusts and cotton emergence relationships. *Transactions of the ASAE*. 25: 1484–1487.
- Bravo-Hollis, H. and Scheinvar, L. 1995. El interesante mundo de las cactáceas. Fondo de Cultura Económica, México, D.F. 233 p.
- Bullock, S.H., Martijena, N.E., Webb, R.H. and Turner, R.M. 2005. Twentieth century demographic changes in cirio and cardón in Baja California, México. *Journal of Biogeography*. 32(1): 127–143. <https://doi.org/10.1111/j.1365-2699.2004.01152.x>
- Celaya-Michel, H. and Castellanos-Villegas, A.E. 2011. Mineralización de nitrógeno en el suelo de zonas áridas y semiáridas. *Terra Latinoamericana*. 29(3): 343–356.
- De Mendiburu, F. 2014. *Agricolae: Statistical Procedures for Agricultural Research*. R Package Version 1.2-0. 2014.
- Degli-Esposti, M.D., Lopes De Siqueira, D., Gomes Pereira, P.R., Alvarez Venegas, V.H., Chamhum Salomão, L.C. and Machado Filho, J.A. 2011. Assessment of nitrogenized nutrition of citrus rootstocks using chlorophyll concentrations in the Leaf. *Journal of Plant Nutrition*. 26(6): 1287–1299. <https://doi.org/10.1081/PLN-120020371>
- Delgado-Fernández, M., Escobar Flores, J.G. and Franklin, K. 2017. El cardón gigante (*Pachycereus pringlei*) y sus interacciones con la fauna en la península de Baja California, México. *Acta Universitaria*. 27(5): 11–18. <https://doi.org/10.15174/au.2017.1274>
- Delgado-Sánchez, P., Yáñez-Espinosa, L., Jiménez-Bremont, J.F., Chapa-Vargas, L. and Flores, J. 2013. Ecophysiological and anatomical mechanisms behind the nurse effect: Which are more important? A multivariate approach for cactus seedlings. *PLOS ONE*. 8(11): e81513. <https://doi.org/10.1371/journal.pone.0081513>
- Doan, T.T., Henry-Des-Tureaux, T., Rumpel, C., Janeau, J.L. and Jouquet, P. 2015. Impact of compost, vermicompost and biochar on soil fertility, maize yield and soil erosion in Northern Vietnam: A three year mesocosm experiment. *Science of the Total Environment*. 514: 147–154. <https://doi.org/10.1016/j.scitotenv.2015.02.005>
- Domínguez, J. 2004. State of the art and new perspectives on vermicomposting research. In: *Earthworm Ecology*. CRC Press, pp 401-424. https://doi.org/10.1007/978-981-19-8080-0_2
- Dumroese, R.K., Jacobs, D.F. and Wilkinson, K.M. 2012. Fases de cultivo: Establecimiento y

- crecimiento rápido. In: Producción de plantas en viveros forestales. Comodoro Rivadavia: Centro de Investigación y Extensión Forestal Andino Patagónico, Argentina, pp 133-142.
- Fenner, M. and Thompson, K. 2005. Seedling establishment. In: The Ecology of Seeds. Cambridge University Press, pp 145-162.
- Flexas, J. and Medrano, H. 2002. Drought-inhibition of photosynthesis in C3 plants: Stomatal and non-stomatal limitations revisited. *Annals of Botany*. 89(2): 183–189. <https://doi.org/10.1093/aob/mcf027>
- Flores, L.S., Pérez López, D. de J., Sangerman-Jarquín, D.M., Rubí Arriaga, M., González Huerta, A. and Ramírez Dávila, J.F. 2022. Densidad de población y fertilización orgánica en habas del centro mexicano. *Revista Mexicana de Ciencias Agrícolas*. 13(2): 317–330. <https://doi.org/10.29312/remexca.v13i2.3131>
- Franco-Salazar, V.A. and Véliz, J.A. 2008. Efectos de la salinidad sobre el crecimiento, acidez titulable y concentración de clorofila en *Opuntia Ficus-Indica* (L) mill. *Revista Multidisciplinaria del Consejo de Investigación de la Universidad de Oriente*. 20(1): 12–17.
- Freire, J.D.L., Dos Santos, M.V.F., Dubeux Júnior, C.B., Bezerra Neto, E., Lira, M.D.A., Da Cunha, M.V., Dos Santos, D.C., De Amorim, S.O., De Mello, A.C.L. 2018. Growth of cactus pear cv. Miúda under different salinity levels and irrigation frequencies. *Anais da Academia Brasileira de Ciências*. 90(4): 3893–3900.
- Godínez-Alvarez, H. and Valiente-Banuet, A. 1998. Germination and early seedling growth of Tehuacan Valley cacti species: the role of soils and seed ingestion by dispersers on seedling growth. *Journal of Arid Environments*. 39(1): 21–31. <https://doi.org/10.1006/jare.1998.0376>
- Goldstein, G., Ortega, J.K., Nerd, A. and Nobel, P.S. 1991. Diel patterns of water potential components for the crassulacean acid metabolism plant *Opuntia ficus-indica* when well-watered or droughted. *Plant Physiology*. 95(1): 274-280. <https://doi.org/10.1104/pp.95.1.274>
- González, J.L. and Chueca Sancho, A. 2010. C4 y CAM: características generales y uso en programas de desarrollo de tierras áridas y semiáridas. Homenaje del doctor Julio López Gorgé. Editorial CSIC-CSIC Press. España, Madrid. 197 p.
- Gonzalez-Cortés, A., Reyes-Valdés, M.H., Robledo-Torres, V., Villarreal-Quintanilla, J.A. and Ramírez-Godina, F. 2018. Pre-germination treatments in four prickly pear cactus (*Opuntia* sp.) species from Northeastern Mexico. *Australian Journal of Crop Science*. 12(10): 1676-168.
- González, F. 2012. Las zonas áridas y semiáridas de México y su vegetación. Instituto Nacional de Ecología. México, D.F. 194 p.
- Granados, D., López, F. and Gama, L. 1998. Adaptaciones y estrategias de las plantas de zonas áridas. *Revista Chapingo Serie Ciencias Forestales y del ambiente*. 4(1): 169–178.
- Graves, S., Piepho, H-P., Selzer, L. and Dorai-Raj, S. 2019. Package “multcompView” Title Visualizations of Paired Comparisons.

- Hsiao, T.C. 1973. Plant responses to water stress. *Annual Review of Plant Biology*. 24:519-570. <https://doi.org/10.1146/annurev.pp.24.060173.002511>
- Kaplan, E.L. and Meier, P. 1958. Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association*. 53(282): 457. <https://doi.org/10.1080/01621459.1958.10501452>
- Loza-Cornejo, S. and Terrazas, T. 2011. Morfo-anatomía de plántulas en especies de Pachycereae: ¿hasta cuándo son plántulas? *Boletín de la Sociedad Botánica de México*. 88: 1–13.
- Loza-Cornejo, S., Terrazas, T., López-Mata, L. and Trejo, C. 2003. Características morfoanatómicas y metabolismo fotosintético en plántulas de *Stenocereus queretaroensis* (Cactaceae): Su significado adaptativo. *Interciencia*. 28(2): 83-89-124.
- Lozano-Isla, F., Benites-Alfaro, O.E. and Pompelli, M.F. 2019. GerminaR: An R package for germination analysis with the interactive web application “GerminaQuant for R”. *Ecological Research*. 34: 339– 346. <https://doi.org/10.1111/1440-1703.1275>
- Maguire, J.D. 1992. Speed of germination-aid in selection and evaluation for seedling emergence and vigor. *Crop Science*. 1962(2): 176–177.
- Mandujano, M.D.C., Golubov, J. and Montaña, C. 1997. Dormancy and endozoochorous dispersal of *Opuntia rastroseeds* in the southern Chihuahuan Desert. *Journal of Arid Environments*. 36(2): 259–266. <https://doi.org/10.1006/jare.1996.0210>
- Mantel, N. and Haenszel, W. 1959. Statistical aspects of the analysis of data from retrospective studies of disease. *Journal of the National Cancer Institute*. 22(4): 719–748. <https://doi.org/10.1093/jnci/22.4.719>
- Marchiol, L., Mondini, C., Leita, L. and Zerbi, G. 1999. Effects of municipal waste leachate on seed germination in soil-compost mixtures. *Restoration Ecology*. 7(2): 155–161. <https://doi.org/10.1046/j.1526-100X.1999.72007.x>
- Martínez-Ramos, M., Arroyo-Cosultchi, G., Mandujano, M.C. and Golubov, J. 2016. Dinámica poblacional de *Mammillaria humboldtii* una cactácea endémica de Hidalgo, México. *Botanical Sciences*. 94(2): 199–208. <https://doi.org/10.17129/botsci.270>
- Meléndez, L., Hernández, A. and Fernández, S. 2006. Efecto de la fertilización foliar y edáfica sobre el crecimiento de plantas de maíz sometidas a exceso de humedad en el suelo. *Bioagro*. 18(2): 107–114.
- Moghaieb, R.E.A., Saneoka, H. and Fujita, K. 2004. Effect of salinity on osmotic adjustment, glycinebetaine accumulation and the betaine aldehyde dehydrogenase gene expression in two halophytic plants, *Salicornia europaea* and *Suaeda maritima*. *Plant Science* 5(166): 1345–1349. <https://doi.org/10.1016/j.plantsci.2004.01.016>
- Montaño, N.M., Ayala, F., Bullock, S., Briones, O., García, F., García, R., Maya, Y., Perroni, Y., Siebe, C., Tapia, Y., Troyo, E. and Yépez E. 2016. Almacenes y flujos de carbono en ecosistemas áridos y semiáridos de México: Síntesis y perspectivas. *Terra Latinoamericana*. 34(1): 39–59.

- Moreno, F.L.P. 2009. Respuesta de las plantas al estrés por déficit hídrico. *Agronomía Colombiana*. 27(2): 179–191.
- Naseer, M.N., Rahman, F.U., Hussain, Z., Khan, I.A., Aslam, M.M., Aslam, A., Waheed, H., Khan, A.U. and Iqbal, S. 2022. Effect of salinity stress on germination, seedling growth, mineral uptake and chlorophyll contents of three cucurbitaceae species. *Brazilian Archives of Biology and Technology*. 65: 2022.
- National Research Council. 2002. Predicting Invasions of Nonindigenous Plants and Plant Pests In: Establishment. National Academies Press, United States, Washington (DC).
- Nieto-Garibay, A., Murillo, A.B., Luna, G.P., Troyo, D.E., García, H., Aguilar, G.M., Holguín, P.R. and Larrinaga, M.J.A. 2021. La composta. Importancia, elaboración y uso agrícola. Trillas. México, D.F. 88 p.
- Niklas, K.J., Molina-Freaner, F., Tinoco-Ojanguren, C. and Paolillo, D.J. 2000. Wood biomechanics and anatomy of *Pachycereus pringlei*. *American Journal of Botany*. 87(4): 469– 481. <https://doi.org/10.2307/2656590>
- Niklas, K.J., Molina-Freaner, F., Tinoco-Ojanguren, C. and Paolillo, D.J. 2002. The biomechanics of *Pachycereus pringlei* root systems. *American Journal of Botany*. 89(1): 12–21. <https://doi.org/10.2307/2656590>
- Nilsen, E.T. and Orcutt, D.M. 2000. Physiology of Plants Under Stress: Soil and Biotic Factors. John Wiley & Sons. USA, New York. 680 p.
- Nobel, P.S. 2006. Parenchyma–chlorenchyma water movement during drought for the hemiepiphytic cactus *Hylocereus undatus*. *Annals of Botany*. 97(3): 469-474. <https://doi.org/10.1093/aob/mcj054>
- Nobel, P.S. 1988. Environmental biology of agaves and cacti. U.K., Cambridge. 270 p.
- Ortiz, M., Silva, H., Silva, P. and Acevedo, E. 2003. Estudio de parámetros hídricos foliares en trigo (*Triticum aestivum* L.) y su uso en selección de genotipos resistentes a sequía. *Revista Chilena de Historia Natural*. 76(2): 219–233. <http://dx.doi.org/10.4067/S0716-078X2003000200008>
- Pérez-Sánchez, R.M., Flores, J., Jurado, E. and González-Salvatierra, C. 2015. Growth and ecophysiology of succulent seedlings under the protection of nurse plants in the Southern Chihuahuan Desert. *Ecosphere*. 6(3): 36. <https://doi.org/10.1890/ES14-00408.1>
- Pimienta-Barrios, E., Hernández, J.Z., Muñoz-Urias, A. and Robles-Murguía, C. 2012. Ecophysiology of young stems (cladodes) of *Opuntia ficus-indica* in wet and dry conditions. *Gayana Botánica*. 69(2): 232–239. <http://dx.doi.org/10.4067/S0717-66432012000200002>
- Puente, M.E. 2004. Poblaciones bacterianas endófitas y del rizoplano de plantas del desierto degradadoras de roca y su efecto sobre el crecimiento del cardón. CIBNOR, México, La Paz. 166 p.

- Ramírez, D.A. 2011. Los objetos nodriza como refugio y fuente de nutrientes: Reflexiones sobre el establecimiento y restauración de cactáceas en zonas áridas de la vertiente occidental de los andes. *Ecología Aplicada*. 10(1–2): 83.
- Ranal, M.A. and Santana, D.G. 2006. How and why to measure the germination process? *Blazilian Journal of Botany*. 29: 1–11. <https://doi.org/10.1590/S0100-84042006000100002>
- Salisbury, F.B. and Ross, C.W. 1992. Plant Physiology. Wadsworth Publishing Co., USA, California. 682 p.
- Salmon, Y., Lintunen, A., Dayet, A., Chan, T., Dewar, R., Vesala, T. and Hölttä, T. 2020. Leaf carbon and water status control stomatal and nonstomatal limitations of photosynthesis in trees. *New Phytologist*. 226(3): 690–703. <https://doi.org/10.1111/nph.16436>
- Sánchez-Soto, B., Reyes-Olivas, Á., García-Moya, E. and Terrazas, T. 2010. Germinación de tres cactáceas que habitan la región costera del noroeste de México. *Interciencia*. 35(4): 299-305.
- Sarria-Perea, E. 2010. Efectos de los cationes calcio y magnesio sobre la germinación de semillas de *Juncus* en condiciones de estrés salino. UPV, España, Gandía. 58 p.
- Schuch, U.K. and Kelly, J.J. 2008. Salinity tolerance of cacti and succulents. Turfgrass, Landscape and Urban IPM Research Summary p-155.
- Strain, H.H. and Svec, W.A. 1966. Extraction, separation, estimation and isolation of the chlorophylls. In: The Chlorophylls. Academic Press, pp 21–66.
- Suzán-Azpiri, H. and Sosa, V.J. 2006. Comparative performance of the giant cardon cactus (*Pachycereus pringlei*) seedlings under two leguminous nurse plant species. *Journal of Arid Environments*. 65(3): 351–362. <https://doi.org/10.1016/j.jaridenv.2005.08.002>
- Taiz, L. and Zeiger, E. 2006. Fisiología Vegetal. Universitat Jaume, España. 1338 p.
- Teixeira Roth, V., Castro Cepero, V., Ceroni Stuva, A. and Eyzaguirre Pérez, R. 2004. Diversidad y densidad de la comunidad de cactáceas en el cerro Umarmata y quebrada Orobél en el valle del río Chillón (Lima) y su relación con los factores edáficos. *Ecología Aplicada*. 3(1–2): 1–8.
- Trujillo G., M.E., Méndez N., J.R.; Hossne G., A.J. and Parra D., F.J. 2010. Efecto de la humedad y compactación de un Ultisol de la sabana del estado Monagas sobre la concentración de clorofila y carotenoides, lavado de electrolitos y contenido relativo de agua en plantas de soya. *Acta Universitaria* 20(3): 18–30. <https://doi.org/10.15174/au.2010.65>
- Turner, R.M. 1990. Long-term vegetation change at a fully protected Sonoran Desert site. *Ecology*. 71(2): 464–477. <https://doi.org/10.2307/1940301>
- Uddin, M.B., Mukul, S.A. and Hossain, M.K. 2012. Effects of Organic Manure on Seedling Growth and Nodulation Capabilities of Five Popular Leguminous Agroforestry Tree Components of Bangladesh. *Journal of Forest and Environmental Science*. 28(4): 212–219. <http://dx.doi.org/10.7747/JFS.2012.28.4.212>

- Valiente-Banuet, A. and Ezcurra, E. 1991. Shade as a cause of the association between the cactus *Neobuxbaumia tetetzo* and the nurse plant *Mimosa luisana* in the Tehuacan Valley, Mexico. *The Journal of Ecology*. 79(4): 961. <https://doi.org/10.2307/2261091>
- Vásquez-Méndez, R., Ventura-Ramos, E.J. and Acosta-Gallegos, J.A. 2011. Habilidad de estimación de los métodos de evapotranspiración para una zona semiárida del centro de México. *Revista Mexicana de Ciencias Agrícolas*. 2(3): 399–415.
- Yamasaki, L.R. and Dillenburg, S. 1999. Measurements of leaf relative water content in *Araucaria angustifolia*. *Revista Brasileira de Fisiologia Vegetal*. 11(2): 69–75.